



Highlighted Student Research

Intraspecific variation in leaf traits facilitates the occurrence of trees at the Amazonia–Cerrado transition

Igor Araújo^{a,*}, Beatriz S. Marimon^{a,b}, Marina C. Scalón^{a,c}, Wesley J.A. Cruz^a, Sophie Fauset^d, Tiffani C.S. Vieira^e, David R. Galbraith^f, Manuel U. Gloor^f^a Programa de Pós-Graduação em Ecologia e Conservação, Universidade do Estado de Mato Grosso, Campus Nova Xavantina, Nova Xavantina, MT, Brazil^b Universidade do Estado de Mato Grosso, Departamento de Ciências Biológicas, Nova Xavantina, Brazil^c Programa de Pós-Graduação em Ecologia e Conservação, Universidade Federal do Paraná, Curitiba, PR, Brazil^d University of Plymouth, School of Geography, Earth and Environmental Sciences, Plymouth, Devon, United Kingdom^e Academic of the degree course in Biological Sciences, Universidade do Estado de Mato Grosso, Campus Nova Xavantina, Nova Xavantina, MT, Brazil^f School of Geography, University of Leeds, Leeds, LS2 9JT, UK

ARTICLE INFO

Editor: Hermann Heilmeyer

Keywords:

Leaf morpho-anatomical traits

Savanna–forest gradient

Microclimate

Ecological strategies

Tropics

Climate change

ABSTRACT

The ability of plant species to adjust key functional traits through intraspecific variation may determine their success in persisting on our planet in the future, especially in unstable habitats, such as the Amazonia–Cerrado transition zone. We assessed intraspecific variation in 12 leaf morphological and anatomical traits for four tree species along a savanna–forest gradient, including rocky *cerrado*, typical *cerrado* and *woodland savanna*. Generally, all evaluated species showed great intraspecific variation. Our findings demonstrate that trees occurring in the *woodland savanna* are potentially more vulnerable to climate change, while in the *cerrado* the individuals presented higher tolerance to water deficit and high temperatures. Trees occurring in open-canopy habitats showed smaller stomata, higher stomata and trichome densities, compared to the same species growing in the *woodland savanna*. In contrast, the individuals in the *woodland savanna* shift leaf traits to increase resource acquisition (e.g. light), showing higher specific leaf area and larger stomata, compared to *cerrado* individuals. We have shown that vegetation-induced shifts in leaf morphological and anatomical traits are a major effect in within-species variability, with consequences for persistence and tolerance of species under future climatic conditions.

1. Introduction

Current and future climate change are among the main threats to ecosystems globally (Yu et al., 2019), and it is expected that their impacts and direct consequences will lead to global losses in biodiversity, carbon stocks and ecosystem services (Allen et al., 2010; Collins et al., 2013; Law, 2014; IPCC, 2019; Araújo et al., 2021). The forecast is for increased mortality risk of species across biomes (Van Mantgem et al., 2007; Phillips et al., 2009; Peng et al., 2011; McDowell, 2018), which are already sensitive to the current climate and may be even more vulnerable in the future (Araújo et al., 2021). Climate has changed enough to exceed the physiological limits to which some plants are adapted (Shaw and Etterson, 2012; Araújo et al., 2021) and led to negative effects that compromise survival and persistence of species over time (Becklin et al., 2016). In this context, even greater concerns

arise regarding the ability of species to adapt quickly to climate change (Burrows et al., 2011; Dawson et al., 2011; Hoffmann and Sgrò, 2011). This is especially so for plants, which are sessile organisms with a long lifespan. Thus, trees will need flexibility during their lifetime to deal with rapid environmental and climatic changes, unrelated to genetic changes, and therefore intraspecific variation and phenotypic plasticity can be decisive for the persistence of trees and thus the risk of extinction due to future climate change (Nicotra et al., 2010; Benito Garzón et al., 2011). In this context, functional traits have advanced our understanding of tree physiology, community structure and ecosystem functioning (Wright et al., 2004; De la Riva et al., 2015; Volf et al., 2016).

Recent studies have integrated the variation of intraspecific characteristics in functional traits to investigate the responses of trees along environmental gradients (Laforest-Lapointe et al., 2014; Siefert et al., 2015). Intraspecific variation allows trees to establish under new

* Corresponding author.

E-mail address: igor.araujo@outlook.com.br (I. Araújo).<https://doi.org/10.1016/j.flora.2021.151829>

Received 10 October 2020; Received in revised form 22 March 2021; Accepted 15 April 2021

Available online 18 April 2021

0367-2530/Crown Copyright © 2021 Published by Elsevier GmbH. All rights reserved.

environmental conditions (Joshi et al., 2001; Byars et al., 2007; Maracahipes et al., 2018). In particular, environmental variation induces changes in the functional traits of trees (Anderson and Gezon, 2015; Lázaro-Nogal et al., 2015; Niinemets, 2016), at morphological (Capuzzo et al., 2012), anatomical (Rossatto and Kolb, 2010, 2012; De Paula et al., 2018) and physiological levels (Rossatto et al., 2010). This scenario highlights the set of functional traits that allow plants to survive in contrasting environments has fundamental consequences for our understanding of the dynamics of vegetation, especially in highly complex and hyperdiverse areas such as savanna–forest transitions (Marimon et al., 2014; Oliveras and Yadvinder, 2016).

The Amazonia–Cerrado transition is a natural laboratory in the open, as it presents a marked gradient (savanna–forest) with high diversity of species and contrasting environmental aspects (Marimon-Junior and Haridasan, 2005; Marimon et al., 2014; Araújo et al., 2021), that allow investigating the relationships between functional traits and the environment, providing a step forward to better understand and predict ecological patterns in a changing environment (Hoffmann et al., 2005; Maracahipes et al., 2018). Savannas have low soil fertility, lower air humidity, high temperatures and light intensity, seasonal droughts and frequent fires (Marimon-Junior and Haridasan, 2005; Reis et al., 2015) and can favour trees with conservative resource use strategies (Pausas et al., 2017; Maracahipes et al., 2018) that tolerate such environmental conditions, reflected in leaf traits such as the high leaf water content to avoid desiccation (Bündchen et al., 2015), long petioles to reduce leaf surface temperature preventing thermal leaf damage (Yates et al., 2010), thick leaves to reduce leaf damage (Dahlgren et al., 2006; Rossatto et al., 2010). In addition, savanna-adapted species can display a set of anatomical traits to prevent excessive water loss such as the high density of trichomes and stomata (Gianoli and Gonzalez-Teuber, 2005; Galmés et al., 2007) and also smaller stomata sizes and maximum opening of the stomata pore that benefits more efficient control in the use of water (Franco, 2002; Rossatto et al., 2009). On the other hand, trees in forests tend to present resource acquisition strategies (Hoffmann et al., 2012), with a combination of functional traits linked to greater uptake of light, water and nutrients. For example, it is expected to find species with high specific leaf area, thin leaves, larger stomata sizes and maximum opening of the stomatal pore (Grime, 1979; Westoby, 1998; Ogburn and Edwards, 2010) that promote higher growth rates.

Although interest in functional traits has grown more and more in plant ecology, in hyperdiverse and complex tropical systems, such as the most diverse transition on the planet between the Amazonia and the Cerrado, we still do not know how the variation in functional traits, especially the anatomical traits (e.g., stomata density, stomata size and maximum opening of the stomatal pore) differ between ecological scales, such as vegetation, species and individuals (Albert et al., 2010; Messier et al., 2010; Salazar et al., 2018). This fact has made it difficult to determine the scale-dependent aspects of the variation in functional traits (Messier et al., 2010). At the individual level, the variability in the functional traits of trees is influenced by climatic drivers (e.g., temperature and precipitation) (Fajardo and Piper, 2011; Richardson et al., 2013). In addition, soil texture, successional stage and disturbance regimes can also contribute to this variation (Salazar et al., 2018). In terms of vegetation, the availability of water and nutrients, soil compaction, salinity and microclimate variations can induce variations in traits (Messier et al., 2010). However, at the species level, the variation may result from interspecific competition for resources such as water, light and nutrients (Schemske et al., 2009; Albert et al., 2010; Messier et al., 2010). As far as we know, no research has used a wide scan of leaf anatomical traits to measure the change in traits on a series of ecological scales. Knowing which scales have the greatest variation in functional traits will provide important explanations about the causes of variability in functional traits and may contribute to subsidies on patterns and processes on spatial and temporal scales that are ecologically more significant (McGill et al., 2006).

In this study we report how the variation in leaf functional traits (e.

g., morphological and anatomical) changes in three nested ecological scales (vegetations, species and individuals) along a savanna–forest gradient in the Amazonia–Cerrado transition, which is making it increasingly dry and hot and where some species can already be affected by climate change (Tiwari et al., 2020; Araújo et al., 2021). In addition, we have introduced little-explored valuable information on the variability of intraspecific characteristics in leaf functional traits of co-occurring trees in contrasting environments in tropical systems. We test two hypotheses: (1) among ecological scales, vegetation will contribute to greater variation in leaf traits, since the distinct habitats present contrasting environmental aspects, as shown in Marimon-Junior and Haridasan (2005) and Figure S1; (2) species have significant intraspecific variability in leaf functional traits irrespective of the habitat. We expected smaller and thicker leaves with higher trichome density in habitats under higher incidence of light, high temperature, and low water availability. On the other hand, larger and thinner leaves with larger stomata will be found in woodland savanna vegetation, a habitat with lower light incidence, lower temperature, and higher water availability (Rossatto et al., 2010; Rossatto and Kolb, 2010, 2012; Capuzzo et al., 2012; Reich et al., 2014; Souza et al., 2015).

2. Materials and methods

2.1. Study area and species description

We carried out the study at the Bacaba Municipal Park (BMP), a conservation unit of approximately 500 ha, in the municipality of Nova Xavantina-MT, Brazil (Fig. 1), located in the transition region between the Amazonia and Cerrado biomes (Marimon-Junior and Haridasan, 2005; Marimon et al., 2014). The climate shows marked seasonality with two well-defined periods, one rainy (October to March) and the other dry (April to September; Alvares et al., 2013), with average annual rainfall of 1.500 mm, elevation of 340 m (Marimon et al., 2010). Mean monthly temperature is 25 °C, ranging from 18.7 to 33.5 °C, according to data obtained from INMET (Instituto Nacional de Meteorologia –<http://www.inmet.gov.br>) considering data from 14 years period (2004–2019).

The typical *cerrado* occupies 153.77 ha of the BMP, is a savanna phytophysognomy composed of low vegetation cover, with trees spaced from each other, and also presents a dense low stratum (Mews et al., 2011; Gomes et al., 2016). Generally, it occurs on deep, well-drained, dystrophic, acidic Latosols and Neosols (Marimon-Junior and Haridasan, 2005). In contrast, the rocky *cerrado* occupies about 25.44 ha of the BMP area, on shallow soils and with rocky outcrops (Litholic Neosols). In this environment, trees are positioned in the gaps between the rocks, where accumulation and decomposition of organic matter and sand deposition resulting from weathering of the rocks occurs (Maracahipes et al., 2011; Gomes et al., 2016). The rocky *cerrado* has a flora with a high degree of endemism (Alves and Kolbek, 2010) and many endangered species (Mendonça and Lins, 2000). The woodland savanna occupies about 162.02 ha of the BMP area, comprising large trees, with thick litter layer (Marimon et al., 1998; Franczak et al., 2011; Reis et al., 2015) and dystrophic soil (Marimon-Junior and Haridasan, 2005).

For our study we selected four co-occurring species (*Hymenaea stigonocarpa* Mart. ex Hayne – Fabaceae, *Qualea parviflora* Mart. – Vochysiaceae, *Vatairea macrocarpa* (Benth.) Ducke – Fabaceae and *Pseudobombax longiflorum* (Mart.) A. Robyns – Malvaceae) in *cerrado* vegetation (rocky *cerrado* and typical *cerrado*) and a woodland savanna in BMP. These species have extensive geographic distributions and belong to three important botanical families of the Cerrado biome (Ratter et al., 2006), in addition to having high importance value indices locally, in terms of density, frequency and relative dominance (Mews et al., 2011; Reis et al., 2015; Gomes et al., 2016). For each species, we haphazardly selected five individuals per vegetation type, totalling 15 individuals per species.




Fig. 1. Neotropical savannas and woodland savanna in the Amazonia–Cerrado transition, Brazil, South America.

2.2. Leaf morphological and anatomical parameters

For each individual of each species, we selected eight leaves, five for morphological characterization and three leaves for anatomical

Table 1
Leaf morphological and anatomical traits evaluated for tropical trees in the Amazonia–Cerrado transition. ¹ Grime, 1979; ² Bündchen et al., 2015; ³ Poorter and Bongers, 2006; ⁴ Turner, 1994; ⁵ Pallardy, 1981; ⁶ Gratani et al., 2006; ⁷ Westoby, 1998; ⁸ Pearce et al., 2006; ⁹ Weyers and Meidner, 1990; ¹⁰ Ehleringer and Björkman, 1978; ¹¹ Abrams et al., 1994; ¹² Rossatto et al., 2009.

Traits	Acronym	Unit	Functional significance
Specific leaf area	SLA	cm ² g ⁻¹	Resource uptake, resource use efficiency, structural defense, and growth strategies ¹ ;
Leaf water mass content	LWMC	mg g ⁻¹	Leaf temperature reduction and protection against desiccation ² ;
Petiole length	PEL	mm	Light capture and heat dissipation ³ ;
Leaf thickness	LT	mm	Resource acquisition and use, and resistance to physical damage ⁴ ;
Adaxial cuticle thickness	ACT	μm	Water conservation ⁵ ;
Adaxial epidermis thickness	AET	μm	Light and heat dissipation ⁶ ;
Palisade parenchyma thickness	PPT	μm	Photosynthetic rates ⁷ ;
Spongy parenchyma thickness	SPT	μm	Efficient gas exchange ⁸ ;
Stomatal density	STO	mm ⁻²	Carbon assimilation and water use efficiency ⁹ ;
Trichome density	TRD	mm ⁻²	Preventing the increase in leaf temperature ¹⁰ ;
Stomatal size	STS	μm	Water use efficiency ¹¹ ;
Maximum opening of the stomatal pore	AMAX	μm ²	Prevent water loss ¹² ;

determinations (Table 1). As a standardization criterion, we collected fully expanded leaves, exposed to full sunlight, and free of pathogens (i. e. leaf standardization protocol). We measured leaf thickness, specific leaf area (ratio between leaf area and leaf dry mass) and leaf water mass content for each species. We determined the thickness of the fresh leaves with an electronic digital micrometer and digitized them with a scanner to determine leaf area and calculated the specific leaf area (Abramoff et al., 2004). We estimated the wet weight of the leaves with a precision balance (± 0.001 g), then placed them in paper bags in an oven at 60 °C, and after 72 h determined the dry weight. The leaf water content was calculated as the ratio between wet and dry weights (Pérez-Harguindeguy et al., 2013).

For anatomical characterization, we took imprints of the leaf surface with high-tech molding silicon (Speedex), as proposed by Weyers and Johansen (1985). Subsequently, we used colorless enamel to make the impression of the mold on slides, which was visualized with an optical microscope, with an attached camera, to view stomata and trichomes. We took stomatal size and stomatal density measurements under 10 × microscopic amplification selecting haphazardly 10 fields per leaf. Microscope images were processed with the ImageJ software (Abramoff et al., 2004).

We calculated stomatal density, for each individual, as the average number of stomata counted in the same fields of view recorded previously, and then we estimated the average stomatal densities, lengths, and widths per species, measuring 25 stomatal complexes from 3 leaves per individual. We measured the length of the guard cell (“L” in μm), the width of the pair of guard cells (“W”, μm), the size of the stomata (“S”, estimated as S = L * W, according to Franks et al., 2009, 2012) and the maximum area of the stomatal pore (“a_{max}”, μm²). We calculated the maximum area of the stomatal pore as a_{max} = α * S, α = 0.12 (Franks and Beerling, 2009). We determined the trichome density (when present) as the average number of trichomes counted in the same fields of view recorded previously and estimated the densities.

For each of the three leaves described above, we took 2 cm × 2 cm sample from the median portion of the leaf blade and used a freehand

cross-section. We fixed the leaf fragment on a half-open petiole (*Cecropia* sp.) and cut it freehand with the help of a razor blade. Afterwards, we stored the samples in Petri dishes and with the aid of a Pasteur pipette we added 3 ml of sodium hypochlorite, and we waited for 5 to 10 min until the samples became translucent. Then, we rinsed three times with distilled water and stained the samples with a solution of 50% methylene blue and 50% safranin. We photographed the slides with a camera attached to an optical microscope, haphazardly selecting 10 fields from each leaf with a $10\times$ magnification objective to assess the thickness of the adaxial cuticle, adaxial epidermis, and palisade and spongy parenchyma (Roeser, 1962).

2.3. Data analysis

Before the analysis, we tested for normality and homoscedasticity of the data using the Shapiro-Wilk and Levene tests (Levene, 1961; Shapiro-Wilk, 1965). To compare leaf traits between species and vegetation we used split-plot ANOVAs, where species and the interaction between

species and vegetation were nested within vegetation. We used the lmer function from nlme package (Pinheiro et al., 2017) and lsmeans (Lenth and Lenth, 2018) and multcomp (Hothorn et al., 2008) packages for the post-hoc analyses. To test whether humidity and air temperature and the photosynthetic photon flux density (PPFD) were different between vegetation types, we performed univariate analysis of variance (ANOVA) individually for each variable. Subsequently, we used Tukey post-hoc tests to identify significant differences between specific vegetation types. We also calculated the coefficient of variation for the functional traits to observe which traits are more variable and how they varied in terms of species and vegetation types (Garnier et al., 2001). We further performed a principal component analysis (PCA) to verify which functional traits were associated with vegetation types and species.

To understand controls on trait variability, we used different groupings of mixed linear models, adjusting the separate models for each trait (Rosas et al., 2019). Vegetation types, species, and individuals were introduced as nested random factors to assess how the variability of functional traits was distributed among these different levels of




Fig. 2. Trait differences of four tree species from the Amazonia–Cerrado transition that co-occur in three contrasting vegetation types. Different lowercase letters denote significant differences (Split-plot ANOVA, $P < 0.05$).

organization. All variables were checked for normality and $\log_{10}$ -transformed whenever necessary to ensure normality. All data analyses were performed using the R program, version 3.6.1 (R Core Team, 2019) with a 5% significance level.

3. Results

3.1. Distribution of leaf morphological and anatomical traits

Qualea parviflora and *V. macrocarpa* showed higher stomatal density and trichome density, regardless of the vegetation type they occurred, while *H. stigonocarpa* and *P. longiflorum* were generally characterized by higher petiole length and leaf water mass content (Fig. 2). The first two axes of the PCA explained 65.1% of the variation and presented a very evident difference between species, which allowed identification of different groups (Fig. 3). PC1 explained 37.4% of the variation and was related to differences between species, while PC2 explained 27.7% of the variation, and separated the vegetation types.

We verified the separation of the three vegetation types (PC2), with a clear distinction between those with extreme environmental conditions such as the rocky *cerrado* (shallow and rocky soil, high temperature and high light intensity) and the woodland *savanna* (deep soil, low light intensity and high humidity), while the typical *cerrado* occupied an intermediate position (Fig. 3). Adaxial cuticle thickness presented the highest correlation with PC2 (Table S1).

3.2. Variability of leaf morphological and anatomical traits at different organizational levels

For most of the traits evaluated, vegetation type and species together explained most of the variance (Fig. 4). The vegetation type explained between 42 and 85% of the variation in adaxial epidermis thickness, adaxial cuticle thickness, and specific leaf area. Species explained between 68 to 91% of the variation in stomatal size, stomatal density, leaf water mass content, and maximum opening of the stomatal pore. On the other hand, some traits showed low variation at the individual level, which varied from 4 to 11% of the variation in adaxial cuticle thickness, leaf water mass content and adaxial epidermis thickness (Fig. 4). To access the decomposition of variance for all organizational levels, see Table S2.

3.3. Intraspecific variability of leaf morphological and anatomical traits

The effect of species was significant for most traits, except for adaxial epidermis thickness (Table S3). The interactions between the two effects were significant, except for leaf water mass content (Table S3). These effects (i.e. species and vegetation types) will be examined successively, combining the results shown in Table S3 with the data obtained in all individual species (Fig. 2).

Patterns within species across vegetation types showed that individuals occurring on *cerrado* formations generally have more conservative strategies than those in *woodland savanna* formations. For example, specific leaf area, a trait associated with resource acquisition, was higher for individuals in the *woodland savanna* ($P < 0.001$; Fig. 2) while leaf water mass content, associated with resistance to desiccation, was higher for individuals of the typical *cerrado* and rocky *cerrado* ($P = 0.010$; Fig. 2).

Similarly, individuals occurring on *cerrado* formations, especially those on the rocky *cerrado*, generally had higher spongy parenchyma thickness, compared to individuals of the same species in the *woodland savanna* ($P < 0.001$; Fig. 2). On the other hand, the trichomes density and the thickness of the cuticle and adaxial epidermis and the palisade parenchyma showed differences with the species and vegetation interaction ($P < 0.001$; Fig. 2).

For other traits, differences across vegetation types were less consistent across species. For example, petiole length widely varied among individuals, being highest for *Q. parviflora* individuals in the rocky *cerrado*, whereas for *H. stigonocarpa* and *P. longiflorum*, they were higher in the *woodland savanna* ($P = 0.003$; Fig. 2). Similarly, stomatal traits did not exhibit consistent variation across vegetation types. For *Q. parviflora* and *V. macrocarpa*, stomatal density was higher for individuals in the *cerrados*, whereas for *P. longiflorum* the stomata density values were higher in the *woodland savanna* ($P = 0.002$; Fig. 2). *Qualea parviflora* recorded the highest stomatal size values in the rocky *cerrado*, whereas *V. macrocarpa* and *P. longiflorum* recorded the highest values in the *woodland savanna* ($P < 0.001$; Fig. 2).

In general, for all traits, the coefficient of variation was variable at the species level (Figure S2): higher values for *P. longiflorum* in the rocky *cerrado*, for *Q. parviflora* in the typical *cerrado* and for *V. macrocarpa* in the *woodland savanna* (Figure S2). *Hymenaea stigonocarpa* generally showed the lowest variability in terms of species and vegetation (Figure S2). The coefficient of variation was always higher for species in the *woodland savanna* and decreased with the canopy opening (Figure S2). For all species, the most variable trait in the rocky *cerrado*




Fig. 3. Principal component analysis of the morphological and anatomical traits of four tree species from the Amazonia–Cerrado transition, co-occurring in rocky *cerrado*, typical *cerrado* and woodland *savanna*. Maximum opening of the stomatal pore (AMAX), stomata size (STS), stomata density (STO), trichome density (TRD), spongy parenchyma thickness (SPT), palisade parenchyma thickness (PPT), adaxial epidermis thickness (AET), adaxial cuticle thickness (ACT), petiole length (PEL), leaf thickness (LT), leaf water mass content (LWMC) and specific leaf area (SLA).




Fig. 4. Partitioning of the variance of the nested linear models of the morphological and anatomical traits of four species in the Amazonia–Cerrado transition. Maximum opening of the stomatal pore (AMAX), stomata size (STS), stomata density (STO), trichome density (TRD), spongy parenchyma thickness (SPT), palisade parenchyma thickness (PPT), adaxial epidermis thickness (AET), adaxial cuticle thickness (ACT), petiole length (PEL), leaf thickness (LT), leaf water mass content (LWMC) and specific leaf area (SLA). Within means the residual error, all data were transformed (log-10) before analysis.

was the spongy parenchyma thickness while palisade parenchyma thickness was the most variable in the typical *cerrado* and leaf thickness was most variable in the *woodland savanna* (Figure S2).

4. Discussion

The species here evaluated in the largest and most diverse transition between biomes on the planet, the Amazonia–Cerrado, showed variability in their leaf morphological and anatomical traits. Spongy parenchyma thickness, palisade parenchyma thickness and leaf thickness were the most variable traits at the intraspecific level. On the other hand, petiole length and adaxial epidermis thickness are the least variable. These variations between the leaf morphological and anatomical traits are driven by different factors such as species, environmental and genetic modification (Garnier et al., 2001; Cássia-Silva et al., 2017; Smedt et al., 2018). Variations in leaf traits diverged widely between organizational levels. The variance explained by vegetation, species, and individuals allowed us to understand separate contribution to leaf traits' variability. In this study, intraspecific variation of the thickness of the cuticle and the adaxial epidermis were mostly explained by vegetation, which may be driven by water availability, nutrients, and microclimate variations (Marimon-Junior and Haridasan, 2005). Surprisingly, vegetation did not influence leaf water content, which >90% of its variability was explained by species, which may be related to interspecific differences in ecological strategies to use and compete for resources (Albert et al., 2010; Messier et al., 2010). The stomatal dimensions (i.e., the maximum opening of the stomatal pore, stomata size and stomata density) exhibited greater variation also explained at the species level, which may be more related to water and nutrient limitations (Marimon-Junior and Haridasan, 2005; Schemske et al., 2009; Albert et al., 2010). On the other hand, a relatively small percentage of variation (4–35%) was explained by the individual level for all leaf traits, which may be due to the marked climate seasonality and edaphic contrasts in this region (Marimon et al., 2014).

Therefore, our findings show that plants co-occurring in open and closed-canopy environments adopt different ecological strategies to establish in vegetation with contrasting environmental properties (Hoffmann and Franco, 2003; Laureto and Cianciaruso, 2015; Araújo et al., 2021). Our results are corroborated by another large-scale study conducted by Maracahipes et al. (2018), who found intraspecific differences in five traits across generalist species occurring in savanna and forest formations in the Cerrado Biome. Here, we provide a new set of little-explored leaf characteristics, including anatomical traits, which can be integrated into the group of key characteristics that describe the ecological strategies of plant species, not only in transition areas but to

understand the responses of vegetation facing climate change worldwide (Gillison, 2019). In addition, we show that, in savanna-forest systems that occur side by side, regardless of the size of the scale, environmental differences drive changes in functional diversity of vegetation.

Variations in the traits of trees that occur in vegetation with contrasting abiotic stresses can be interpreted as a result of the pressure of disruptive selection (Lemos-Filho et al., 2008) which reflects the functional divergence between individuals of the same species that are adapted to different conditions (Hoffmann and Franco, 2008). Therefore, intraspecific variation found for the species evaluated here can provide a role in adapting to environmental and climatic changes. It also contributes to expanding the ecological and geographical distribution of the species (Bedetti et al., 2011). We showed that, within species, traits associated with drought tolerance (such as cuticle thickness and trichome density) showed higher values in habitats with greater canopy opening, while traits related to productivity (e.g. specific leaf area) were higher in closed-canopy habitats. In addition, species with high variation in traits usually occupy broader ecological niche and can occur in different habitats (Jung et al., 2010).

There was a clear distinction between *woodland savanna* and other vegetation types, mainly driven by differences in leaf morphology and anatomy, reflected in the principal component analysis (Fig. 3). While *woodland savanna* individuals showed higher specific leaf area compared to individuals of the same species occurring in the other *cerrado* formations, rocky *cerrado* individuals showed thicker cuticles, epidermis, palisade and spongy parenchyma reflecting in higher leaf thickness. High specific leaf area is associated with lower investment in the biosynthesis of tissues that provide support and protection (Grime, 1979; Franco et al., 2005). On the other hand, low specific leaf area for individuals growing in *cerrado* is essential to reduce leaf surface area and prevent water loss through transpiration (Poorter et al., 2009). The lower investment in specific leaf area may be a strategy to increase resource retention and allocation to other organs of the plant (Reich et al., 1992; Westoby, 1998; Pérez-Harguindeguy et al., 2013). In addition, the greater investment for the formation of protective tissues and structures, such as the cuticle and epidermis, with thicker cell walls (Fahn and Cutler, 1992), helps to reduce the harmful effects of ultraviolet radiation and to prevent leaf damage caused by herbivores and high temperatures (Turner, 1994; Rozendaal et al., 2006). Under high temperature, vapor pressure deficit and light leaf mesophyll tissues can be affected (Grime, 1979; Reich et al., 1999; Dahlgren et al., 2006; Gratani et al., 2006; Rossatto and Kolb, 2010), resulting in leaf area and tissue thickness changes (Goulet and Bellefleur, 1986) as observed for the evaluated species (Figure S3).

In the *cerrado* formations, individuals also showed higher trichome density and higher stomata density for *Q. parviflora* and *V. macrocarpa*. These traits represent an adaptive strategy to prevent or control the increase in leaf temperature (Ehleringer and Björkman, 1978), as well as reducing water loss (Gianoli and Gonzalez-Teuber, 2005). On the other hand, in *P. longiflorum* stomata density was higher for individuals in the *woodland savanna*, which represents greater CO₂ assimilation and, consequently, higher evapotranspiration rates that promote greater growth of the species (Ogburn and Edwards, 2010). Stomata size and petiole length differences varied depending on the species considered. *Vatairea macrocarpa* and *P. longiflorum* showed smaller stomata in individuals from *cerrado* formations. Smaller stomata and stomatal openings allow more efficient control over water loss (Abrams et al., 1994; Franco, 2002; Goldstein et al., 2008), especially in areas under high temperatures, intense light and low humidity (Shields, 1950; Ackerly, 2004; Poorter et al., 2009), as the studied *cerrados*. *Hymenaea stigonocarpa* and *P. longiflorum* showed longer petioles in individuals located in the *woodland savanna* and shorter petioles for those from *cerrado* formations. In general, petioles with greater length allow greater light uptake (Poorter and Bongers, 2006), which is an advantageous trait in *woodland savanna* environments, where species are subject to greater competition for light (Carswell et al., 2000). Moreover, longer petioles also contribute to reducing leaf clumping and overlapping around the stem (Takenaka, 1994; Poorter, 2009), which can be a key feature in *woodland savanna*. On the other hand, *Q. parviflora* showed longer petioles in individuals that grow in the rocky *cerrado*, which could probably contribute to reduce leaf surface temperature, dissipating heat and preventing thermal leaf damage (Yates et al., 2010; Ye et al., 2011; Leigh et al., 2012).

Together, these characteristics are important to increase the efficiency of plants in the conservation and use of water (Pallardy, 1981; Rossatto and Kolb, 2010), dissipate excessive light and heat (Feller, 1996; Gratani et al., 2006) and promote an increase in photosynthetic rates (Pearce et al., 2006). In fact, the greater water availability in the *woodland savanna* soil (Marimon-Junior and Haridasan, 2005) may allow individuals to invest in traits not only related to water storage and drought tolerance (Larcher, 2003; Monteiro et al., 2016; Lin et al., 2017), while in *cerrado* formations these strategies could be crucial for the individual survival.

Future climate changes are predicted to increase extreme weather events, such as droughts and heatwaves (Meehl and Tebaldi, 2004; Collins et al., 2013). These changes may negatively affect the ecophysiological fitness of the trees and, consequently, increase the risk of mortality, especially for species in the Amazonia–Cerrado transition that already experience critical levels of temperature increase (Araújo et al., 2021), about 2-fold greater than the average heating recorded for the planet (Coe et al., 2016; Hoegh-Guldberg et al., 2018). Therefore, intraspecific variation may represent a key mechanism for species survival in future climates. In this case, functional traits that allow greater water storage and water use efficiency, and assist in the dissipation of heat may be more advantageous for trees subject to extreme weather events (Nicotra et al., 2010; Benito Garzón et al., 2011).

The high intraspecific variation observed for the species in this study is important for persistence over time, especially in this region, where climate change is already affecting trees (Tiwarei et al., 2020; Araújo et al., 2021). Therefore, we suggest that these species are good candidates for the recovery of degraded areas in the Amazonia–Cerrado transition. This aspect is of ultimate importance, since this region is known as the “Arc of deforestation” due to seasonal droughts, fires and uncontrolled deforestation occurring concurrently (Schmidt and Eloy, 2020; Brando et al., 2020). Moreover, the Amazonia–Cerrado transition, which suffers more deforestation than forests and savannas in each individual biome, is close to collapse due to the intense use of land for crops and pastures (Marques et al., 2020). This scenario is particularly concerning because this transitional region comprises greater local biodiversity than each biome separately (Marimon et al., 2006; Mews

et al., 2012; Marimon et al., 2014).

Adding up to previous studies across distinct environmental gradients (Scalon et al., 2017; Silva et al., 2019; Xu et al., 2020), we showed that intraspecific variability in leaf traits is substantial and should not be ignored in trait-based studies, especially in heterogeneous environments, such as transitional zones (Albert et al., 2010; Jung et al., 2010). Our findings showed that widespread species in the Amazonia–Cerrado transition have different functional strategies to establish and maintain populations in contrasting environments through their high potential for intraspecific variation in their functional traits. Whereas trees in the *cerrado* have leaf morphological and anatomical traits linked to competition for water and conservation of resources, trees of the same species growing in *woodland savanna* have traits that increase efficiency in the competition for light and generate higher productivity. Our results highlight the importance of the interaction between the environment and the leaf morphological and anatomical traits for understanding how selective pressures affect the functional diversity of species in tropical ecosystems.

CRediT authorship contribution statement

Igor Araújo: Conceptualization, Formal analysis, Investigation, Methodology, Writing - original draft. **Beatriz S. Marimon:** Conceptualization, Methodology, Writing - review & editing. **Marina C. Scalon:** Conceptualization, Methodology, Writing - review & editing. **Wesley J. A. Cruz:** Formal analysis. **Sophie Fauset:** Writing - review & editing. **Tiffani C.S. Vieira:** Methodology, Investigation. **David R. Galbraith:** Formal analysis, Writing - review & editing. **Manuel U. Gloor:** Formal analysis, Writing - review & editing.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We are thank CAPES (Coordination for the Improvement of Higher Education Personnel) - Finance Code 001 for granting the author's scholarship.

References

- Abramoff, M.D., Magalhães, P.J., Ram, S.J., 2004. Image processing with ImageJ. *Biophotonics International* 11, 36–42.
- Abrams, M.D., Kubiske, M.E., Mostoller, S.A., 1994. Relating wet and dry year ecophysiology to leaf structure in contrasting temperate tree species. *Ecology* 75, 123–133.
- Ackerly, D., 2004. Functional strategies of chaparral shrubs in relation to seasonal water deficit and disturbance. *Ecological Monographs* 74, 25–44.
- Albert, C.H., Thuiller, W., Yoccoz, N.G., Soudant, A., Boucher, F., Saccone, P., Lavorel, S., 2010. Intraspecific functional variability: extent, structure and sources of variation. *Journal of Ecology* 98, 604–613.
- Allen, C.D., Macalady, A.K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., Hogg, E.T., 2010. A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management* 259, 660–684.
- Alvares, C.A., Stape, J.L., Sentelhas, P.C., de Moraes, G., Leonardo, J., Sparovek, G., 2013. Köppen's climate classification map for Brazil. *Meteorologische Zeitschrift* 22, 711–728.
- Alves, R.J.V., Kolbek, J., 2010. Can campo rupestre vegetation be floristically delimited based on vascular plant genera? *Plant Ecology* 207, 67–79.
- Anderson, J.T., Gezon, Z.J., 2015. Plasticity in functional traits in the context of climate change: a case study of the subalpine forb *Boechera stricta* (Brassicaceae). *Global Change Biology* 21, 1689–1703.
- Araújo, I., Marimon, B.S., Scalon, M.C., Fauset, S., Junior, B.H.M., Tiwarei, R., Gloor, M. U., 2021. Trees at the Amazonia–Cerrado transition are approaching high temperature thresholds. *Environmental Research Letters* 16, 1–8.
- Becklin, K.M., Anderson, J.T., Gerhart, L.M., Wadgymar, S.M., Wessinger, C.A., Ward, J. K., 2016. Examining plant physiological responses to climate change through an evolutionary lens. *Plant Physiology* 172, 635–649.

- Bedetti, C.S., Aguiar, D.B., Jannuzzi, M.C., Moura, M.Z., Silveira, F.A., 2011. Abiotic factors modulate phenotypic plasticity in an apomictic shrub [*Miconia albicans* (SW.) Triana] along a soil fertility gradient in a Neotropical savanna. *Australian Journal of Botany* 59, 274–282.
- Benito Garzón, M., Alfa, R., Robson, T.M., Zavala, M.A., 2011. Intra-specific variability and plasticity influence potential tree species distributions under climate change. *Global Ecology and Biogeography* 20, 766–778.
- Bündchen, M., Boeger, M.R.T., Reissmann, C.B., 2015. Estrutura foliar de espécies lenhosas de dossel e sub-bosque em uma floresta subtropical do sul do Brasil. *Iheringia. Série Botânica* 70, 105–114.
- Burrows, M.T., Schoeman, D.S., Buckley, L.B., Moore, P., Poloczanska, E.S., Brander, K. M., Halpern, B.S., 2011. The pace of shifting climate in marine and terrestrial ecosystems. *Science* 334, 652–655.
- Brando, P., Macedo, M., Silvério, D., Rattis, L., Paolucci, L., Alencar, A., Amorim, C., 2020. Amazon wildfires: Scenes from a foreseeable disaster. *Flora* 268, 1–5.
- Capuzzo, J.P., Rossatto, D.R., Franco, A.C., 2012. Differences in morphological and physiological leaf characteristics between *Tabebuia aurea* and *T. impetiginosa* is related to their typical habitats of occurrence. *Acta Botanica Brasiliica* 26, 519–526.
- Carswell, F.E., Meir, P., Wandelli, E.V., Bonates, L.C.M., Kruijt, B., Barbosa, E.M., Jarvis, P.G., 2000. Photosynthetic capacity in a central Amazonian rain forest. *Tree Physiology* 20, 179–186.
- Cássia-Silva, C., Cianciaruso, M., Maracahipes, L., Collevatti, R., 2017. When the same is not the same: phenotypic variation reveals different plant ecological strategies within species occurring in distinct Neotropical savanna habitats. *Plant Ecology* 218, 1221–1231.
- Coe, M.T., Silverio, D.V., Bustamante, M., Macedo, M., Shimbo, J., Brando, P.M., 2016. Feedbacks between land cover and climate changes in the Brazilian Amazon and Cerrado biomes. Paper presented at the AGU Fall Meeting Abstracts.
- Collins, M., Knutti, R., Arblaster, J., Dufresne, J.L., Fichet, T., Friedlingstein, P., Krinner, G., 2013. Long-term climate change: projections, commitments and irreversibility. In: Stocker, T.F et al. (Eds.), *Climate Change 2013 – The Physical Science Basis: Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, pp. 1029–1136. Cambridge University Press.
- Dahlgren, J.P., Eriksson, O., Bolmgren, K., Strindell, M., Ehrlén, J., 2006. Specific leaf area as a superior predictor of changes in field layer abundance during forest succession. *Journal of Vegetation Science* 17, 577–582.
- Dawson, T.P., Jackson, S.T., House, J.I., Prentice, I.C., Mace, G.M., 2011. Beyond predictions: biodiversity conservation in a changing climate. *Science* 332, 53–58.
- De Paula, L.F.A., Kolb, R.M., Porembski, S., Silveira, F.A.O., Rossatto, D.R., 2018. Rocks and leaves: Can anatomical leaf traits reflect environmental heterogeneity in inselberg vegetation? *Flora* 250, 91–98.
- Ehrlinger, J.R., Björkman, O., 1978. Pubescence and leaf spectral characteristics in a desert shrub, *Encelia farinosa*. *Oecologia* 36, 151–162.
- Fahn, A., Cutler, D.F., 1992. Xerophytes. Gebrüder Borntraeger. Stuttgart, Germany.
- Feller, L.C., 1996. Effects of nutrient enrichment on leaf anatomy of dwarf *Rhizophora mangle* L. (red mangrove). *Biotropica* 28, 13–22.
- Franco, A.C., 2002. Ecophysiology of woody plants. In: Marquis, R.J. (Ed.), *The Cerrados of Brazil: Ecology and Natural History of a Neotropical Savanna*, pp. 178–197. Columbia University Press, New York.
- Franco, A.C., Bustamante, M., Caldas, L.S., Goldstein, G., Meinzer, F.C., Kozovits, A.R., Coradin, V.T.R., 2005. Leaf functional traits of Neotropical savanna trees in relation to seasonal water deficit. *Trees* 19, 326–335.
- Franzczak, D.D., Marimon, B.S., Hur Marimon-Junior, B., Mews, H.A., Maracahipes, L., Oliveira, E.A., 2011. Changes in the structure of a savanna forest over a six-year period in the Amazon–Cerrado transition, Mato Grosso state. *Brazil. Rodriguésia* 62, 425–436.
- Franks, P.J., Beerling, D.J., 2009. Maximum leaf conductance driven by CO₂ effects on stomatal size and density over geologic time. *Proceedings of the National Academy of Sciences* 106, 10343–10347.
- Franks, P.J., Drake, P.L., Beerling, D.J., 2009. Plasticity in maximum stomatal conductance constrained by negative correlation between stomatal size and density: an analysis using *Eucalyptus globulus*. *Plant, Cell and Environment* 32, 1737–1748.
- Galmés, J., Flexas, J., Savé, R., Medrano, H., 2007. Water relations and stomatal characteristics of Mediterranean plants with different growth forms and leaf habits: responses to water stress and recovery. *Plant and Soil* 290, 139–155.
- Garnier, E., Laurent, G., Bellmann, A., Debain, S., Berthelot, P., Ducout, B., Navas, M.L., 2001. Consistency of species ranking based on functional leaf traits. *New Phytologist* 152, 69–83.
- Gianoli, E., González-Teuber, M., 2005. Environmental heterogeneity and population differentiation in plasticity to drought in *Convolvulus chilensis* (Convolvulaceae). *Evolutionary Ecology* 19, 603–613.
- Gillison, A.N., 2019. Plant functional indicators of vegetation response to climate change, past present and future: II. Modal plant functional types as response indicators for present and future climates. *Flora* 254, 31–58.
- Goldstein, G., Meinzer, F.C., Bucci, S.J., Scholz, F.G., Franco, A.C., Hoffmann, W.A., 2008. Water economy of Neotropical savanna trees: six paradigms revisited. *Tree Physiology* 28, 395–404.
- Gomes, L., Maracahipes, L., Reis, S.M., Marimon, B.S., Marimon-Junior, B.H., Lenza, E., 2016. Dynamics of the woody vegetation of two areas of Cerrado sensu stricto located on different substrates. *Rodriguésia* 67, 859–870.
- Goulet, F., Bellefleur, P., 1986. Leaf morphology plasticity in response to light environment in deciduous tree species and its implication on forest succession. *Canadian Journal of Forest Research* 16, 1192–1195.
- Gratani, L., Covone, F., Larcher, W., 2006. Leaf plasticity in response to light of three evergreen species of the Mediterranean maquis. *Trees* 20, 549–558.
- Grime, J.P., 1979. *Plant Strategies and Vegetation Processes*. John Wiley & Sons, Chichester.
- Hothorn, T., Bretz, F., Westfall, P., 2008. Simultaneous inference in general parametric models. *Biometrical Journal: Journal of Mathematical Methods in Biosciences* 50, 346–363.
- Hoegh-Guldberg, O., Jacob, D., Taylor, M., Bindi, M., Brown, S., Camilloni, I., Engelbrecht, F., 2018. Impacts of 1.5 °C Global Warming on Natural and Human Systems. *IPCC Special Report*. <http://hdl.handle.net/10138/311749>.
- Hoffmann, A.A., Sgro, C.M., 2011. Climate change and evolutionary adaptation. *Nature* 470, 479–485.
- Hoffmann, W.A., Franco, A.C., 2008. The importance of evolutionary history in studies of plant physiological ecology: examples from cerrados and forests of central Brazil. *Brazilian Journal of Plant Physiology* 20, 247–256.
- Hoffmann, W.A., Geiger, E.L., Gotsch, S.G., Rossatto, D.R., Silva, L.C.R., Lau, O.L., Franco, A.C., 2012. Ecological thresholds at the savanna–forest boundary: how plant traits, resources and fire govern the distribution of tropical biomes. *Ecology Letters* 15, 759–768.
- IPCC, 2019. *Land Degradation: IPCC Special Report on Climate Change, Desertification, Land 5 Degradation, Sustainable Land Management, Food Security, and 6 Greenhouse Gas Fluxes in Terrestrial Ecosystems*. *Sciences* 1–43.
- Jung, V., Violle, C., Mondy, C., Hoffmann, L., Muller, S., 2010. Intraspecific variability and trait-based community assembly. *Journal of Ecology* 98, 1134–1140.
- Larcher, W., 2003. *Physiological Plant Ecology: Ecophysiology and Stress Physiology of Functional Groups*. Springer Science & Business Media.
- Law, B.E., 2014. Regional analysis of drought and heat impacts on forests: current and future science directions. *Global Change Biology* 20, 3595–3599.
- Lázaro-Nogal, A., Matesanz, S., Godoy, A., Pérez-Trautman, F., Gianoli, E., Valladares, F., 2015. Environmental heterogeneity leads to higher plasticity in dry-edge populations of a semi-arid Chilean shrub: insights into climate change responses. *Journal of Ecology* 103, 338–350.
- Leigh, A., Sevanto, S., Ball, M.C., Close, J.D., Ellsworth, D.S., Knight, C.A., Vogel, S., 2012. Do thick leaves avoid thermal damage in critically low wind speeds? *New Phytologist* 194, 477–487.
- Lemos-Filho, J.P., Goulart, M.F., Lovato, M.B., 2008. Populational approach in ecophysiological studies: the case of *Plathymenia reticulata*, a tree from Cerrado and Atlantic Forest. *Brazilian Journal of Plant Physiology* 20, 205–216.
- Lenth, R., Lenth, M.R., 2018. Package ‘lsmmeans’. *The American Statistician* 34, 216–221.
- Levene, H., 1961. Robust tests for equality of variances. *Contributions to probability and statistics. Essays in honor of Harold Hotelling* 279–292.
- Lin, H., Chen, Y., Zhang, H., Fu, P., Fan, Z., 2017. Stronger cooling effects of transpiration and leaf physical traits of plants from a hot dry habitat than from a hot wet habitat. *Functional Ecology* 31, 2202–2211.
- Maracahipes, L., Lenza, E., Marimon, B.S., de Oliveira, E.A., Pinto, J.R.R., Junior, B.H.M., 2011. Estrutura e composição florística da vegetação lenhosa em cerrado rupestre na transição Cerrado-Floresta Amazônica, Mato Grosso. *Brasil. Biota Neotropica* 11, 1–9.
- Maracahipes, L., Carlucci, M.B., Lenza, E., Marimon, B.S., Marimon Jr, B.H., Guimarães, F.A., Cianciaruso, M.V., 2018. How to live in contrasting habitats? Acquisitive and conservative strategies emerge at inter- and intraspecific levels in savanna and forest woody plants. *Perspectives in Plant Ecology* 34, 17–25. *Evolution and Systematics*.
- Marimon, B.S., Felfili, J.M., Lima, E.S., Duarte, W.M.G., Marimon-Junior, B.H., 2010. Environmental determinants for natural regeneration of gallery forest at the Cerrado/Amazonia boundaries in Brazil. *Acta Amazonica* 40, 107–118.
- Marimon, B.S., Marimon-Junior, B.H., Feldpausch, T.R., Oliveira-Santos, C., Mews, H.A., Lopez-Gonzalez, G., Maracahipes, L., 2014. Disequilibrium and hyperdynamic tree turnover at the forest–cerrado transition zone in southern Amazonia. *Plant Ecology and Diversity* 7, 281–292.
- Marimon, B.S., Varella, R.F., Marimon-Junior, B.H., 1998. Fitossociologia de uma área de cerrado de encosta em Nova Xavantina, Mato Grosso. *Boletim do Herbário Ezechias Paulo Heringer* 3, 1–12.
- Marimon, B.S., Lima, E.S., Duarte, T.G., Chiaregatto, L.C., Ratter, J.A., 2006. Observations on the vegetation of northeastern Mato Grosso, Brazil. IV. An analysis of the Cerrado–Amazonian Forest ecotone. *Edinburgh Journal of Botany* 63, 323–341.
- Marimon-Junior, B.H., Haridasan, M., 2005. Comparação da vegetação arbórea e características edáficas de um cerrado e um cerrado sensu stricto em áreas adjacentes sobre solo distrófico no leste de Mato Grosso. *Brasil. Acta Botanica Brasiliica* 19, 913–926.
- Marques, E.Q., Marimon-Junior, B.H., Marimon, B.S., Matricardi, E.A.T., Mews, H.A., Colli, G.R., 2020. Redefining the Cerrado–Amazonia transition: implications for conservation. *Biodiversity and Conservation* 29, 1–17.
- McDowell, N.G., 2018. Deriving pattern from complexity in the processes underlying tropical forest drought impacts. *New Phytologist* 219, 841–844.
- McGill, B.J., Enquist, B.J., Weiher, E., Westoby, M., 2006. Rebuilding community ecology from functional traits. *Trends in Ecology & Evolution* 21, 178–185.
- Meehl, G.A., Tebaldi, C., 2004. More intense, more frequent, and longer lasting heat waves in the 21st century. *Science* 305, 994–997.
- Mendonça, M.P., Lins, L.V., 2000. Lista vermelha das espécies ameaçadas de extinção da flora. Fundação Biodiversitas, de Minas Gerais.
- Messier, J., McGill, B.J., Lechowicz, M.J., 2010. How do traits vary across ecological scales? A case for trait-based ecology. *Ecology Letters* 13, 838–848.
- Mews, H.A., Marimon, B.S., Maracahipes, L., Franzcak, D.D., Marimon-Junior, B.H., 2011. Dinâmica da comunidade lenhosa de um Cerrado Típico na região Nordeste do Estado de Mato Grosso. *Brasil. Biota Neotropica* 11, 1–10.

- Mews, H.A., Marimon, B.S., Ratter, J.A., 2012. Observations on the vegetation of Mato Grosso, Brazil. V. Changes in the woody species diversity of a forest in the Cerrado–Amazonian Forest Transition zone and notes on the forests of the region. *Edinburgh Journal of Botany* 69, 239–253.
- Monteiro, M.V., Blauša, T., Verhoef, A., Hadley, P., Cameron, R.W.F., 2016. Relative importance of transpiration rate and leaf morphological traits for the regulation of leaf temperature. *Australian Journal of Botany* 64, 32–44.
- Nicotra, A.B., Atkin, O.K., Bonser, S.P., Davidson, A.M., Finnegan, E.J., Mathesius, U., Valladares, F., 2010. Plant phenotypic plasticity in a changing climate. *Trends in Plant Science* 15, 684–692.
- Niinemets, Ü., 2016. Leaf age dependent changes in within-canopy variation in leaf functional traits: a meta-analysis. *Journal of Plant Research* 129, 313–338.
- Ogburn, R.M., Edwards, E.J., 2010. The ecological water-use strategies of succulent plants. *Advances in Botanical Research* 55, 179–225.
- Pallardy, S.G., 1981. Closely related woody plants. In: Kozlowski, T.T. (Ed.), *Water Deficits and Plant Growth*. Academic Press, New York, pp. 511–548.
- Pausas, J.G., Keeley, J.E., Schwillk, D.W., 2017. Flammability as an ecological and evolutionary driver. *Journal of Ecology* 105, 289–297.
- Pearce, D.W., Millard, S., Bray, D.F., Rood, S.B., 2006. Stomatal characteristics of riparian poplar species in a semi-arid environment. *Tree Physiology* 26, 211–218.
- Peng, C., Ma, Z., Lei, X., Zhu, Q., Chen, H., Wang, W., Zhou, X., 2011. A drought-induced pervasive increase in tree mortality across Canada's boreal forests. *Nature Climate Change* 1, 467–471.
- Pérez-Harguindeguy, N., Diaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., Gurvich, D.E., 2013. Corrigendum to: new handbook for standardised measurement of plant functional traits worldwide. *Australian Journal of Botany* 64, 715–716.
- Phillips, O.L., Aragão, L.O.C., Lewis, S.L., Fisher, J.B., Lloyd, J., López-González, G., Quesada, C.A., 2009. Drought sensitivity of the Amazon rainforest. *Science* 323, 1344–1347.
- Poorter, H., Niinemets, Ü., Poorter, L., Wright, I.J., Villar, R., 2009. Causes and consequences of variation in leaf mass per area (LMA): a meta-analysis. *New Phytologist* 182, 565–588.
- Poorter, L., 2009. Leaf traits show different relationships with shade tolerance in moist versus dry tropical forests. *New Phytologist* 181, 890–900.
- Poorter, L., Bongers, F., 2006. Leaf traits are good predictors of plant performance across 53 rain forest species. *Ecology* 87, 1733–1743.
- Core Team, R., 2019. R Foundation for Statistical Computing. Vienna, Austria.
- Ratter, J.A., Bridgewater, S., Ribeiro, J.F., 2006. Biodiversity patterns of the woody vegetation of the Brazilian Cerrado. In: Pennington, R.T., Ratter, J.A. (Eds.), *Neotropical Savannas and Seasonally Dry Forests*. CRC Press, pp. 46–81.
- Reich, P.B., 2014. The world-wide 'fast-slow' plant economics spectrum: a traits manifesto. *Journal of Ecology* 102, 275–301.
- Reich, P.B., Ellsworth, D.S., Walters, M.B., Vose, J.M., Gresham, C., Volin, J.C., Bowman, W.D., 1999. Generality of leaf trait relationships: a test across six biomes. *Ecology* 80, 1955–1969.
- Reich, P.B., Walters, M.B., Ellsworth, D.S., 1992. Leaf life-span in relation to leaf, plant, and stand characteristics among diverse ecosystems. *Ecological Monographs* 62, 365–392.
- Reis, S.M., Lenza, E., Marimon, B.S., Gomes, L., Forsthofer, M., Morandi, P.S., Elias, F., 2015. Post-fire dynamics of the woody vegetation of a savanna forest (Cerradão) in the Cerrado–Amazon transition zone. *Acta Botanica Brasiliica* 29, 408–416.
- Roeser, K.R., 1962. Die Nadel der Schwarzkiefer – Massenprodukt und Kunstwert der Naturen. *Mikrokosmos* 61, 31–36.
- Rosas, T., Mencuccini, M., Barba, J., Cochard, H., Saura-Mas, S., Martínez-Vilalta, J., 2019. Adjustments and coordination of hydraulic, leaf and stem traits along a water availability gradient. *New Phytologist* 223, 632–646.
- Rossatto, D.R., Hoffmann, W.A., Franco, A.C., 2009. Características estomáticas de pares congeneros de cerrado e mata de galeria crescendo numa região transicional no Brasil Central. *Acta Botanica Brasiliica* 23, 499–508.
- Rossatto, D.R., Kolb, R.M., 2010. *Gochnatia polymorpha* (Less.) Cabrera (Asteraceae) changes in leaf structure due to differences in light and edaphic conditions. *Acta Botanica Brasiliica* 24, 605–612.
- Rossatto, D.R., Takahashi, F.S.C., Silva, L.C.R., Franco, A.C., 2010. Características funcionais de folhas de sol e sombra de espécies arbóreas em uma mata de galeria no Distrito Federal. Brasil. *Acta Botanica Brasiliica* 24, 640–647.
- Rozendaal, D.M.A., Hurtado, V.H., Poorter, L., 2006. Plasticity in leaf traits of 38 tropical tree species in response to light; relationships with light demand and adult stature. *Functional Ecology* 20, 207–216.
- Salazar, P.C., Navarro-Cerrillo, R.M., Cruz, G., Villar, R., 2018. Intraspecific leaf functional trait variability of eight *Prosopis pallida* tree populations along a climatic gradient of the dry forests of northern Peru. *Journal of Arid Environments* 152, 12–20.
- Scalon, M.C., Haridasan, M., Franco, A.C., 2017. Influence of long-term nutrient manipulation on specific leaf area and leaf nutrient concentrations in savanna woody species of contrasting leaf phenologies. *Plant and Soil* 421, 233–244.
- Schmidt, I.B., Eloy, L., 2020. Fire regime in the Brazilian Savanna: Recent changes, policy and management. *Flora* 268, 1–5.
- Shapiro, S.S., Wilk, M.B., 1965. An analysis of variance test for normality (complete samples). *Biometrika* 52, 591–611.
- Shaw, R.G., Etterson, J.R., 2012. Rapid climate change and the rate of adaptation: insight from experimental quantitative genetics. *New Phytologist* 195, 752–765.
- Shields, L.M., 1950. Leaf xeromorphy as related to physiological and structural influences. *The Botanical Review* 16, 399–447.
- Silva, M.C., Teodoro, G.S., Bragion, E.F., Van Den Berg, E., 2019. The role of intraspecific trait variation in the occupation of sharp forest–savanna ecotones. *Flora* 253, 35–42.
- Smedt, P., Ottaviani, G., Wardell-Johnson, G., Sýkora, K.V., Mucina, L., 2018. Habitat heterogeneity promotes intraspecific trait variability of shrub species in Australian granite inselbergs. *Folia Geobotanica* 53, 133–145.
- Souza, M.C., Franco, A.C., Haridasan, M., Rossatto, D.R., Araújo, J.F., Morellato, L.P.C., Habermann, G., 2015. The length of the dry season may be associated with leaf scleromorphism in cerrado plants. *Anais da Academia Brasileira de Ciências* 87, 1691–1699.
- Takenaka, A., 1994. Effects of leaf blade narrowness and petiole length on the light capture efficiency of a shoot. *Ecological Research* 9, 109–114.
- Tiwari, R., Gloor, E., da Cruz, W.J.A., et al., 2020. Photosynthetic quantum efficiency in south-eastern Amazonian trees may be already affected by climate change. *Plant, Cell and Environment* 48, 1–12.
- Turner, I.M., 1994. Sclerophylly: primarily protective? *Functional Ecology* 8, 669–675.
- Van Mantgem, P.J.J.P., Stephenson, N.L., 2007. Apparent climatically induced increase of tree mortality rates in a temperate forest. *Ecology Letters* 10, 909–916.
- Volf, M., Redmond, C., Albert, Á.J., Le Bagousse-Pinguet, Y., Biella, P., Götzenberger, L., de Bello, F., 2016. Effects of long- and short-term management on the functional structure of meadows through species turnover and intraspecific trait variability. *Oecologia* 180, 941–950.
- Westoby, M., 1998. A leaf–height–seed (LHS) plant ecology strategy scheme. *Plant and Soil* 199, 213–227.
- Weyers, J.D.B., Johansen, L.G., 1985. Accurate estimation of stomatal aperture from silicone rubber impressions. *New Phytologist* 101, 109–115.
- Weyers, J.D.B., Meidner, H., 1990. *Methods in Stomatal Research*. Longman, Harlow.
- Wright, I.J., Reich, P.B., Westoby, M., Ackerly, D.D., Baruch, Z., Bongers, F., Villar, R., 2004. The worldwide leaf economics spectrum. *Nature* 428, 821–827.
- Xu, W., Tomlinson, K.W., Li, J., 2020. Strong intraspecific trait variation in a tropical dominant tree species along an elevational gradient. *Plant Diversity* 42, 1–6.
- Yates, M.J., Verboom, G.A., Rebelo, A.G., Cramer, M.D., 2010. Ecophysiological significance of leaf size variation in Proteaceae from the Cape Floristic Region. *Functional Ecology* 24, 485–492.
- Ye, C., Li, M., Hu, J., Cheng, Q., Jiang, L., Song, Y., 2011. Highly reflective superhydrophobic white coating inspired by poplar leaf hairs toward an effective “cool roof. *Energy and Environmental Science* 4, 3364–3367.
- Yu, D., Liu, Y., Shi, P., Wu, J., 2019. Projecting impacts of climate change on global terrestrial ecoregions. *Ecological Indicators* 103, 114–123.